



Cannabis Oil against the benefits they expect you will experience.

If you have any concerns or queries about using this medicine, please speak to your doctor or pharmacist. Keep this leaflet with your medicine. You may need to reference it in the future.

OzSun 20:1 oil

Delta-9-tetrahydrocannabinol (THC) 20mg/mL,
Cannabidiol (CBD) <1mg/mL
Medicinal Cannabis Oil - For oral use

Consumer Medicine Information

This leaflet addresses frequently asked questions about OzSun Cannabis Oils. This leaflet does not contain all available information and does not replace speaking to your trusted doctor or pharmacist.

To ensure you have the most up-to-date information on these products, OzSun recommends utilising the advice and experience of your doctor and pharmacist.

All medicines present potential risks and benefits. You are receiving this leaflet because your doctor has weighed the risks of you using OzSun

Precautions And Warnings

OzSun 20:1 Oil is contraindicated (should not be used) for:

1. Women who are pregnant or lactating;
2. Persons with a known or suspected cannabinoid allergy;
3. Persons with serious cardiovascular disease e.g., congestive heart failure; and
4. Persons with personal history of schizophrenia or psychosis.

OzSun 20:1 Oil should be used with caution for:

1. Persons with suspected or confirmed family history of serious psychiatric disorders such as schizophrenia, psychosis, major depression, bipolar disorder or personality disorder; and
2. Persons with a history of alcohol or drug abuse or substance dependence.

Do not take OzSun 20:1 Oil after the expiry date has passed.

Do not take OzSun 20:1 Oil if the packaging is damaged or shows signs of tampering.

Dependence on OzSun 20:1 Oil is unlikely but may occur in some persons.

Psychoactive side effects attributed to THC may occur in some persons. It is recommended to avoid consuming alcohol while consuming OzSun 20:1 Oil.

Before you start using OzSun 20:1 Oil, inform your doctor if:

1. You have allergies to any other medicines;
2. You have allergies to cannabinoids or terpenes;
3. You are pregnant or are planning on becoming pregnant;
4. You have serious liver or kidney problems;
5. You have experienced serious heart problems such as angina, previous heart attacks, uncontrolled or poorly controlled high blood pressure or an irregularity with your heart rate or heartbeat;
6. You are taking any blood thinners.
7. You are elderly, have difficulty performing activities of daily living (ADLs) such as preparing food or drink, or are at risk to falls and instability; or
8. You are taking any other medicines, including prescription, OTC, and herbal medicines.

If you have not informed your doctor about any of the above, tell them before you begin using OzSun 20:1 Oil.

Dosage

OzSun 20:1 Oil is for oral use only. Do not use OzSun 20:1 Oil without dosing and administration advice from your doctor. If you are unsure about administration or dosage, please consult with your prescribing doctor or pharmacist. Currently, there is no standardised or clinically defined dosage for OzSun 20:1 Oil. Following the 'start low, go slow' philosophy of cannabis medicine, all patients will be started on a low dose and titrated under clinical supervision until a therapeutic response is achieved or side effects limit titration. Your optimal dose will be established by you and your doctor by monitoring the effectiveness of each dose and observing for any side effects. Due to the long half-life of oral cannabis medicines, it is recommended to wait a minimum of 6-8 hours between doses. Titration should be conducted in a slow and mindful manner in small increments with the guidance of your prescribing doctor.

Each bottle of OzSun 20:1 Oil is supplied with a reusable oral dosing syringe to ensure simple administration

Administration

1. Remove child resistant bottle cap by firmly pushing down on the cap, squeeze and rotate counter clockwise. Tamper proof seal will detach on first opening.
2. Insert adapter (inside the cap) into bottle until it is in a flush position.
3. Hold bottle in vertical position and firmly push dosing syringe into the adapter in bottle, ensuring syringe is secure in small central adapter hole.
4. Turn bottle upside down, slowly pull back on plunger, drawing oil into the syringe from bottle.
5. Pull plunger to desired dose as represented on syringe in printed increments. Carefully check dose.
6. **Turn bottle right way up before removing syringe from fitting to ensure oil does not leak.**
7. Place tip of syringe inside the mouth, under the tongue and slowly push down plunger.
8. Replace child resistant bottle cap firmly pushing down, squeeze and rotate clockwise.
9. After every use, wash with warm water and return bottle and syringe

If You Forget A Dose

If you forget to take a dose, take the next dose when you remember and continue to take the next dose at the regular time and interval. If you forget a dose, do not consume a double dose to make up for it as this may increase the risk of experiencing unwanted side effects. If you are unsure or need guidance, please call your doctor or pharmacist.

If You Take Too Much

Common side effects of a THC overdose include anxiety and paranoia. Whilst this may be unpleasant, the effects are generally short lasting.

If you have preexisting cardiovascular disease, chest pain or shortness of breath please call 000. If you are experiencing suicidal thoughts and ideations, please call lifeline or 000. Once the episode has subsided, please reduce your dose and talk to your prescribing doctor about dose adjustments or medication changes.

Safety Information

OzSun 20:1 Oil contains the psychoactive cannabinoid THC. THC ingestion can result in temporary intoxication, confusion, fatigue, drowsiness and impairment. It is strongly advised that patients refrain from driving or operating heavy machinery while on THC-based medicines. It is currently an offence in Australia to drive with a detectable amount of THC in the body. For more information on legalities surrounding cannabis medicines, please speak to your doctor or pharmacist.

THC Predominant Medications

THC is a powerful medicine, even at very low doses. It is also associated with most cannabis side effects. It is intoxicating and impairing at high doses and frequent overuse of THC based medications can down-regulate the body's own endocannabinoid system (ECS). THC based medications need to be titrated in a mindful and reflective manner. It is best to start in the sub-therapeutic range and slowly titrate up to therapeutic doses over 1-2 weeks.

Side Effects Of THC

(generally related to high doses - safe and well tolerated when used appropriately)

- Asthenia (abnormal physical weakness or lack of energy)
- Confusion
- Disorientation
- Dizziness
- Drowsiness
- Vertigo
- Balance and coordination problems
- Dry mouth
- Hallucinations
- Anxiety
- Paranoid thoughts
- Increased appetite
- Vomiting or nausea
- Short term memory impairment
- Light-headedness
- Uncontrolled laughter or euphoria
- Low mood

Storage

All medicines (including cannabis medicines) must be stored safely in a location that children and animals cannot reach. Light and heat can degrade cannabis products. Store all cannabis products in a cool dark place below 25 degrees celsius, away from direct sunlight.

Do not transfer product material to a different container.

Do not use the product after the expiry date printed on the label.

Sponsor Details

Sponsored & manufactured by
ECS Botanics MC Pty Ltd.
info@ecs-botanics.com

This information was prepared on
21 January 2025

WARNING:
THIS MEDICINAL CANNABIS
PRODUCT IS A SCHEDULE 8
POISON: PRESCRIPTION ONLY
MEDICINE KEEP OUT OF REACH
OF CHILDREN

If you have any other questions about medicinal cannabis products, please talk to your Medical Practitioner.